



Diet after a Nissen Fundoplication

After a Nissen fundoplication, you may experience difficulty swallowing due to swelling at the site of the surgery.

After Nissen Fundoplication surgery, your diet will change slowly from clear liquids to full liquids to a soft diet and then progress to a regular diet.

A soft diet is recommended for the first 2 to 6 weeks following surgery. You will then slowly advance to a regular diet; depending on your progress and tolerance for food.

The foods tolerable on this diet can vary greatly from one person to the next.

To speed your recovery, we recommend you follow a soft diet, and follow these suggestions:

- Eat small, frequent meals to prevent your stomach from stretching. We recommend eating 6 times per day, or every 2 to 3 hours.
- Eat foods that are easy to swallow and digest. These usually consist of soft, moist foods such as soup, gelatin, pudding, and yogurt. Avoid gummy foods such as bread and tough meats.
- Cut all foods in small pieces (the size of your pinky fingernail)
- Take small bites, chew your food well, and avoid gulping. This helps prevent gas build-up and aids in swallowing.
- To prevent swallowing air, which produces excess gas, avoid drinking through a straw and don't chew gum or tobacco.
- Avoid caffeine, carbonated drinks, alcohol, citrus & tomato products.
- Avoid any foods that you know cause stomach gas and distention, including corn, beans, peas, lentils, onions, broccoli, cauliflower, and cabbage.
- You may find it helpful to use a blender to puree foods
- Remember to chew foods well before swallowing.

The Soft Diet

A soft diet serves as a transition from liquids to a regular diet. It generally eliminates foods that are difficult to chew or swallow and also spicy, fried, or gas-producing foods.

To ease digestion, the following foods are generally eliminated:

- Tough meats
- Raw fruits and vegetables
- Chewy or crispy breads
- Nuts and seeds
- Fried, greasy foods
- Spicy foods
- High fiber foods like whole grain bread

Food Group	Foods Recommended	Foods to Avoid
Beverages	Milk Decaf tea Powdered drink mixes Non-citrus juices	Chocolate drinks Caffeinated drinks Coffee Carbonated drinks Alcohol Citrus juices
Grains	Ready to eat cereal Refined bread Crackers Plain white rice Pasta Pancakes/waffles	Whole-grain breads and cereals Anything with seeds, nuts, raisins, dried fruit, or coconut Whole grain rice Sweet rolls, coffee cake, or donuts Popcorn & potato chips
Proteins Meats and beans	Tender meat, pork, poultry, and fish (baked, broiled, boiled, roasted, stewed, or simmered) Eggs Tofu Smooth peanut butter	Anything fried Tough meats with gristle Sausage Shellfish Fatty meats Cold cuts/lunch meat Nuts and seeds Crunchy peanut butter
Dairy	Low-fat milk products Smooth yogurt Mild cheese Cottage cheese	Sharp/strong cheeses Dairy with nuts or seeds Cheese with peppers

Vegetables	Soft-cooked or canned vegetables Fresh lettuce or tomato Potatoes without the skin Peeled white or sweet potatoes (boiled, mashed, baked or creamed)	Raw vegetables, tomatoes, Gas-producing vegetables: broccoli, Brussels sprouts, cabbage, onions, cauliflower, corn, green peppers, cucumber, radishes, sauerkraut
Fruit	Soft raw fruits (without skin) Cooked and canned fruits Fruit Juice	Dried fruits Fruits with skins, seeds or pits, such as berries, figs or raisins All citrus fruits and juices Ripe bananas Coconut
Fats and Sweets	Ice cream sherbet and frozen yogurt Pudding Cake and cookies without hard pieces Sugar, syrup, honey, jelly, Seedless jam Molasses Marshmallows Butter and margarine Mayonnaise and vegetable oils Mildly seasoned salad dressings, sauces, and gravies Plain cream cheese and sour cream	Spicy salad dressings Bacon, bacon fat, ham fat, lard, salt pork Fried foods Nuts Anything with dried fruit, nuts, coconut, candied fruit Peanut brittle

If you have any additional questions after your surgery,
please feel free to call our office at 724-439-1020
and ask to speak to a medical assistant.